

Health Disparities in Perinatal Mental Health Resource Document

Health Disparities in Perinatal Mental Health Workgroup: Collaboration of the Women Psychiatrists Caucus, Black Psychiatrists Caucus, Council on Women's Mental Health, Council on Minority Mental Health and Health Disparities

Workgroup Members:

Garima Arora, M.D.
Uju Berry, M.D.
Helen Blaisdell-Brennan, M.D.
Nancy Byatt, M.D.
Anjali Gupta, M.D. (Chair)
Carla Hammond, M.D.
Jacqueline A. Hobbs,
M.D.,Ph.D
Pamela Montano, M.D.
Jennifer Payne, M.D.
Laika Simeon-Thompson, M.D.
Catherine Wei, M.D.
Carole Warshaw, M.D.
Rochelle Woods, M.D.
Muhammad Zeshan, M.D.

Introduction

APA's Health Disparities in Perinatal Mental Health Resource Document — developed by the APA's Health Disparities in Perinatal Mental Health Work Group — is an evolving resource for members who want to learn more about health disparities in women's perinatal mental health.

Health disparities exist in perinatal maternal mental health. Black and Hispanic women have higher rates of postpartum depression and are at increased risk of developing perinatal anxiety disorders. Women of color also experience higher rates of maternal suicide and substance use disorders during the perinatal period. Lower socioeconomic status, lack of insurance, geographic location, age, having a dominant language other than English, being single, and experiencing abuse by an intimate partner, also increase the risk of perinatal mental health issues for women.

Understanding health disparities, training in culturally competent practices, enhancing screening and support services, including for IPV, trauma, and SDOH, and providing psychoeducation to decrease stigma is important at the provider level. Additionally at the systemic level, reducing barriers to care, improving access to culturally competent care, and advocating for policies that reduce social and political determinants of health are all ways to continue to address health disparities for women and their mental health

during the perinatal period

Included below are relevant APA documents as well as education and resources to better understand and address health disparities.

Access

While there are many evidence-based practices for perinatal mood and anxiety disorders, screening rates vary widely, and $\leq 25\%$ of those who screen positive for depression receive any treatment. Gaps in care loom largest for individuals who are Black/African American, Hispanic/Latina/o/e/x, American Indian or Alaska Native, or Native Hawaiian or other Pacific Islander (populations that have been marginalized).

While effective treatments are available, care often falls short of patients' needs and of recommended standards. Barriers persist across the mental healthcare pathway (screening, assessment, and treatment until symptom remission), including stigma, fear regarding treatment, and a dearth of community-based mental health supports. Providers and patients report that barriers are amplified by 1) gaps in provider training, 2) lack of established treatment workflows, 3) lack of access to mental healthcare resources, referrals, and consultation, 4) lack of integrated responses to IPV and other lifetime trauma (vs. under SDOH?), and 5) microaggressions, implicit racial biases, and discrimination. Gaps and inequities persist across the perinatal mental healthcare pathway. Thus, the US lags behind other high-income countries.

Adverse social determinants of health (SDOH) and disparities in screening or treatment rates drive inequities in perinatal mental healthcare. SDOH are the social circumstances in which people are born, work, live, and age and the wider set of forces shaping daily life. SDOH influence 80% of health outcomes as well as patient engagement with the healthcare system. Addressing patients' adverse SDOH is critical to improving health equity and is essential for high-quality healthcare. Professional organizations and policy makers recommend screening for adverse SDOH at healthcare visits. Integrating social services into mental healthcare is critical for improving health equity and patient experience. While healthcare and community providers want to address inequities in access to care, they do not have the training, skills, resources, or tools to do so. This dearth of resources, combined with racial discrimination in healthcare and community settings, contribute to 1) higher rates of depression and anxiety symptoms and 2) lower screening and treatment rates among marginalized populations.

Postpartum Depression and Anxiety

Women of color are disproportionately less likely to receive accurate diagnosis and treatment for perinatal mood and anxiety disorders than white women. Limited access to healthcare contributes to this lack of treatment. Additionally, cultural stigmas and taboos surrounding mental health in specific communities contribute to underreporting and underdiagnosis of perinatal mood and anxiety disorders. This disparity is also exacerbated by bias in the healthcare system. Ensuring we continue to develop culturally-sensitive screening tools and treatment plans and train a healthcare workforce that is culturally aware, compassionate, and knowledgeable of the ways that SDOH, IPV, and trauma across the lifespan affect diagnosis and treatment is important.

Postpartum Psychosis

Racial disparities in postpartum psychosis reflect inequities in the diagnosis, treatment, and outcomes for women of color. Postpartum psychosis is a rare but severe condition that can occur after childbirth, requiring prompt intervention due to its potential for life-threatening consequences.

Black and Hispanic women often face delayed diagnoses or are misdiagnosed with other conditions such as depression or schizophrenia, rather than postpartum psychosis. There are a number of contributing factors. Stereotypes and biases in mental health care contribute to these disparities. Symptoms in women of color may be dismissed or underestimated, leading to a failure in timely recognition of the severity of their condition. Socioeconomic disparities play a significant role. Women of color tend to have less access to high-quality perinatal mental health care due to systemic factors such as lack of insurance, geographic barriers, and limited access to culturally competent providers. Cultural beliefs around motherhood and mental health may also create barriers to openly discussing mental health concerns. Mental health issues, including postpartum psychosis, may be stigmatized in some racial and ethnic communities. This stigma can discourage women from seeking help or sharing their symptoms, worsening the risk of untreated psychosis. Additionally, healthcare providers may harbor implicit biases which can lead them to underestimate the severity of symptoms and the necessity of prompt intervention that may result in less aggressive or comprehensive interventions compared to white women. Delays in diagnosis and treatment can lead to severe consequences, including suicide, infanticide, or long-term psychiatric illness.

Research on postpartum psychosis has focused on predominantly white populations, limiting the understanding of how this condition manifests and impacts women of different racial and ethnic backgrounds. This lack of data contributes to the challenges in creating effective, targeted interventions.

Mortality Rates

Maternal mortality refers to deaths due to complications from pregnancy or up to a year following childbirth. Suicide, drug overdose, and homicide are currently the leading causes of pregnancy-related deaths in the United States. There are a number of disparities that impact maternal mortality including limited access to quality care, race/ethnicity, low-income, lack of education, and language barriers.

Black women are disproportionately affected, with maternal mortality rates approximately three times higher than those of white women. This disparity persists regardless of income or education level, highlighting systemic issues in healthcare access and quality. Indigenous women also experience high maternal mortality rates, nearly twice as high as white women. While the maternal mortality rate for Latina women is lower than that for black and indigenous women, it is still higher than that for white women. Asian and Pacific Islander women tend to have lower maternal mortality rates compared to other racial minorities, closer to those of white women. However, disparities can exist within subgroups and based on geographic location. White women have the lowest maternal mortality rates among racial groups, but these rates have been rising over time.

Contributing factors include limited access to healthcare services. Poverty, lack of education, IPV, and inadequate insurance coverage disproportionately affect minorities, impacting maternal health and mental health outcomes. Implicit biases in healthcare and historical mistrust contribute to these disparities. Racial and ethnic minorities often face barriers to accessing quality prenatal and maternal healthcare. Conditions like hypertension, diabetes, and obesity, which are more prevalent in some racial groups, increase maternal mortality risks. Improving healthcare access, addressing social determinants of health, and increasing awareness and training among healthcare providers to deliver culturally competent care are all ways to reduce these disparities.

Substance Use Disorders

Lower socioeconomic status contributes to financial stressors, unstable housing situations, and food insecurities which can increase the use of substances as a coping mechanism. Pregnant persons of color (POC) are at increased risk of substance use disorders (SUD) during pregnancy that can lead to complications during pregnancy and birth. Lower social support can also contribute to substance use during pregnancy. Limited access to prenatal care and SUD treatment can further impact risk of morbidity and mortality.

Systemic racism intersects with other forms of discrimination and results in increased chronic stress which can make pregnant persons of color more vulnerable to substance misuse. This effect can be exacerbated when it intersects with social determinants of health such as poverty and IPV which can independently increase the risk of substance use disorders.

There is limited availability for specialized SUD programs designed for pregnant persons. Additionally, lack of access to SUD treatment disproportionately affects those with limited financial resources. Intra and extra community stigma related to substance use against minority perinatal patients can create barriers to care. Systemic racism also makes it more likely that a pregnant person of color would face harsher penalties for substance use compared with white women. This may be a barrier for pregnant persons of color to seek help. Other barriers include limited access to culturally and linguistically appropriate care and affordable access to care.

Trauma and Intimate Partner Violence (IPV)

It is important to understand the links between IPV, maternal mental health, and SUDs when assessing for these conditions. Experiencing IPV has a significant impact on maternal mental health and substance use and is a major contributor to maternal mortality, suicide, overdose, and homicide. People who experience IPV often face additional barriers to care by abusive partners who actively undermine their mental health, coerce them to use substances, prevent them from accessing treatment, control their medications, sabotage their recovery efforts, and then leverage stigma-related policies against them, particularly in relation to child custody loss, as part of a broader pattern of abuse and control (Phillips et al., 2024; Warshaw et al., 2014). These policies disproportionately impact women from marginalized/Black, Latin, and AI/AN communities and serve as a major deterrent to care.

Relevant APA Documents

[APA Position Statement on Medicaid Coverage for Maternal Postpartum Care](#)

[Position Statement on Screening and Treatment of Mood and Anxiety Disorders During Pregnancy and Postpartum](#)

[Perinatal Mental Health Toolkit](#)

References

Admon LK, Dalton VK, Kolenic GE, Ettner SL, Tilea A, Haffajee RL, Brownlee RM, Zochowski MK, Tabb KM, Muzik M, Zivin K. Trends in suicidality 1 year before and after

birth among commercially insured childbearing individuals in the United States, 2006-2017. *JAMA Psychiatry*. 2021;78(2):171-176.

Alley DE, Asomugha CN, Conway PH, Sanghavi DM. Accountable health communities: addressing social needs through Medicare and Medicaid. *N Engl J Med*. 2016;374(1):8-11.

Arend J, Tsang-Quinn J, Levine C, Thomas D. The patient-centered medical home: history, components, and review of the evidence. *Mt Sinai J Med*. 2012;79(4):433-450.

Berwick DM. A user's manual for the IOM's 'Quality Chasm' report. *Health Aff (Millwood)*. 2002;21(3):80-90.

Bodnar-Deren S, Benn EKT, Balbierz A, Howell EA. Stigma and postpartum depression treatment acceptability among Black and white women in the first six-months postpartum. *Matern Child Health J*. 2017;21(7):1457-1468.

Brown C, Daniel R, Addo N, Knight S. The experiences of medical students, residents, fellows, and attendings in the emergency department: implicit bias to microaggressions. *AEM Educ Train*. 2021;5(S1):S49-S56.

Byatt N, Levin LL, Ziedonis D, Moore Simas TA, Allison J. Enhancing participation in depression care in outpatient perinatal care settings: a systematic review. *Obstet Gynecol*. 2015 Oct;126(5):1048-1058.

Byatt N, Biebel K, Friedman L, Debordes-Jackson G, Ziedonis D, Pbert L. Patient's views on depression care in obstetric settings: how do they compare to the views of perinatal health care professionals? *Gen Hosp Psychiatry*. 2013;35(6):598-604.

Byatt N, Biebel K, Debordes-Jackson G, Lundquist RS, Moore Simas TA, Weinreb L, Ziedonis D. Community mental health provider reluctance to provide pharmacotherapy may be a barrier to addressing perinatal depression: a preliminary study. *Psychiatr Q*. 2013 Jun;84(2):169-174.

Byatt N, Biebel K, Lundquist R, Moore Simas TA, Debordes-Jackson G, Allison J, Ziedonis D. Patient, provider, and system-level barriers and facilitators to addressing perinatal depression. *J Reprod Infant Psychol*. 2012 Dec;30(5):436- 439.

Byatt N, Simas TA, Lundquist RS, Johnson JV, Ziedonis DM. Strategies for improving perinatal depression treatment in North American outpatient obstetric settings. *J Psychosom Obstet Gynaecol*. 2012 Dec;33(4):143-161.

Byatt N, Xu W, Levin LL, Moore Simas TA. Perinatal depression care pathway for obstetric settings. *Int Rev Psychiatry*. 2019;31(3):210-228.

Byhoff E, Freund KM, Garg A. Accelerating the implementation of social determinants of health interventions in internal medicine. *J Gen Intern Med*. 2018;33(2):223-225.

Bodnar-Deren S, Benn EKT, Balbierz A, Howell EA. Stigma and postpartum depression treatment acceptability among Black and white women in the first six-months postpartum. *Matern Child Health J*. 2017;21(7):1457-1468

Campbell, J., Matoff-Stepp, S., Velez, M. L., Cox, H. H., & Laughon, K. Pregnancy-associated deaths from homicide, suicide, and drug overdose: Review of research and the intersection with intimate partner violence. *Journal of Women's Health*, 2021;30(2): 236–244. <https://doi.org/10.1089/jwh.2020.8875>

Clare CA, Yeh J. Postpartum depression in special populations: a review. *Obstet Gynecol Surv*. 2012;67(5):313- 323.

Council on Community Pediatrics. Poverty and Child Health in the United States. *Pediatrics*. 2016;137(4).

Council on Patient Safety in Women's Health Care, American College of Obstetricians and Gynecologists. Patient Safety Bundle: Reduction of Peripartum Racial/Ethnic Disparities. 2016.
<https://safehealthcareforeverywoman.org/wp-content/uploads/Reduction-of-Peripartum-Disparities-Bundle.pdf>.

Cox EQ, Sowa NA, Meltzer-Brody SE, Gaynes BN. The perinatal depression treatment cascade: baby steps toward improving outcomes. *J Clin Psychiatry*. 2016;77(9):1189-1200.

Erving CL, Thomas CS, Frazier C. Is the black-white mental health paradox consistent across gender and psychiatric disorders? *Am J Epidemiol*. 2019;188(2):314-322.

Gavin AR, Melville JL, Rue T, Guo Y, Dina KT, Katon WJ. Racial differences in the prevalence of antenatal depression. *Gen Hosp Psychiatry*. 2011;33(2):87-93.

Gavin AR, Tabb KM, Melville JL, Guo Y, Katon W. Prevalence and correlates of suicidal ideation during pregnancy. *Arch Womens Ment Health*. 2011;14(3):239-246.

Goff SL, Moran MJ, Szegda K, Fioroni T, DeBanate MA, Byatt N. Development and pilot testing of an adaptable protocol to address postpartum depression in pediatric practices serving lower-income and racial/ethnic minority families: contextual considerations. *Implement Sci Commun*. 2020;1:66

Howell EA, Mora PA, Horowitz CR, Leventhal H. Racial and ethnic differences in factors associated with early postpartum depressive symptoms. *Obstet Gynecol*. 2005;105(6):1442-1450.

Hughes LS, Likumahuwa-Ackman S. Acting on social determinants of health: a primer for family physicians. *Am Fam Physician*. 2017;95(11):695-696.

Jackson JS, Torres M, Caldwell CH, Neighbors HW, Nesse RM, Taylor RJ, Trierweiler SJ, Williams DR. The National Survey of American Life: A study of racial, ethnic and cultural influences on mental disorders and mental health. *Int J Methods Psychiatr Res*. 2004;13(4):196-207.

Marmot M. Social determinants of health inequalities. *Lancet*. 2005;365(9464):1099-1104.

National Academy of Sciences. Integrating Social Care into the Delivery of Health Care: Moving Upstream to Improve the Nation's Health. 2019.

National Academies of Sciences, Engineering, and Medicine. Implementing High-Quality Primary Care: Rebuilding the Foundation of Health Care. Washington, DC: The National Academies Press; 2021.

National Institutes of Health. Screening and referral for social determinants of health: Innovative health care applications and future directions. Accessed June 15, 2020 <https://www.scgcorp.com/SocialDeterminants18/>.

National Perinatal Association Position Statement 2017. <https://ncperinatalassociation.org/wp-content/uploads/2019/09/NPA-Bulletin-2017.pdf>

Phillips H, Lyon E, Krans EE, Warshaw C, Chang JC, Pallatino C. Barriers to help-seeking among intimate partner violence survivors with opioid use disorder. *Int Rev Psychiatry*. 2021 Sep;33(6):534-542.

Sandoval RS, Afolabi T, Said J, Dunleavy S, Chatterjee A, Ölveczky D. Building a tool kit for medical and dental students: addressing microaggressions and discrimination on the wards. *MedEdPORTAL*. 2020;16:10893-10893.

Sidebottom A, Vacquier M, LaRusso E, Erickson D, Hardeman R. Perinatal depression screening practices in a large health system: identifying current state and assessing opportunities to provide more equitable care. *Arch Womens Ment Health*. 2021 Feb;24(1):133-144.

Substance Use and Substance Use Disorder Among Pregnant and Postpartum People. <https://www.asam.org/advocacy/public-policy-statements/details/public-policy-statements>

/2022/10/12/substance-use-and-substance-use-disorder-among-pregnant-and-postpartum-people

Substance Use and Substance Use Disorders by Race and Ethnicity, 2015-2018 Issue Brief

<https://aspe.hhs.gov/sites/default/files/documents/784266c8155778feca25050ab9d50996/substance-use-sud-race-ethnicity-2015-2019.pdf>

Tabb KM, Hsieh W-J, Gavin AR, Eigbake M, Faisal-Cury A, Hajaraih SKM, Huang W-hD, Laurent H, Carter D, Nidey N, Ryckman K, Zivin K. Racial differences in immediate postpartum depression and suicidal ideation among women in a Midwestern delivery hospital. *J Affect Disord Rep.* 2020;1:100008.

U.S. DHHS Task Force on Maternal Mental Health. *National Strategy to Improve Maternal Mental Health Care.* 2024

Torres MB, Salles A, Cochran A. Recognizing and reacting to microaggressions in medicine and surgery. *JAMA Surg.* 2019;154(9):868-872.

Warshaw C., Tinnon E., *Coercion Related to Mental Health and Substance Use in the Context of Intimate Partner Violence: A Toolkit for Screening, Assessment, and Brief Counseling in Primary Care and Behavioral Health Settings.* National Center on Domestic Violence, Trauma & Mental Health. Chicago, IL. 2018.

Wang PS, Lane M, Olfson M, Pincus HA, Wells KB, Kessler RC. Twelve-month use of mental health services in the United States: results from the National Comorbidity Survey Replication. *Arch Gen Psychiatry.* 2005;62(6):629-640.

VIGNETTE #1

Visit 1

Isabella, a 24-year-old Spanish-speaking woman with a past history of one previous miscarriage, who is currently 9 weeks pregnant, presents to your psychiatric office after being referred from her PCP for scoring 16 on the GAD-7 at her last PCP visit two weeks ago. While in the waiting room today, she completed the GAD-7 again and scored 18. You speak with her with the aid of a virtual Spanish interpreter. After you introduce yourself and your role, Isabella tells you that she just recently found out about her pregnancy. She says that she has been constantly worried about her baby, her own health, and her future and that she feels restless, cannot sleep at night, and is tired throughout the day. She mentions that she has been getting into more arguments with her partner as well. She is fidgeting in her seat and tapping her foot. She does not make good eye contact with you. She asks if there is a way to help control her worry.

You acknowledge that it is very normal to feel nervous during pregnancy and inquire about any depressive symptoms she may be experiencing, including low mood, decreased interest in activities she usually enjoys, and hopelessness. She denies feeling any symptoms of depression and again states that she has felt increasingly on-edge over the past two months. She has no thoughts of hurting herself, her partner,

or her baby. She denies any substance use and says that her partner has been “as supportive as he can be” while working two jobs. You provide education on various methods that can reduce anxiety, including medication, therapy, and coping skills/techniques. She says that she is not interested in any psychiatric medication at this time, and she requests time to think about whether she wants to start therapy. You take five minutes to share breathing exercises, relaxation techniques, and other tips including exercise, meditation, and journaling that she can try. You provide her with written handouts of these techniques in Spanish that she can take home with her. She agrees to see you again in two weeks to track her progress. Based on your assessment, you believe that her symptoms are consistent with adjustment disorder with anxiety. You recognize that cultural stigmas surrounding mental health in specific communities can lead to underreporting and underdiagnosis of mood disorders among women of color.

Visit 2

Before your next appointment with Isabella, you conduct a brief review of her chart to see if there have been any major medical events since her last appointment. You see a note from an emergency department visit 5 days ago in which she complained of severe nausea and vomiting. She received IV fluids and antiemetic therapy and was discharged home the same day. You look over the labs and see that she had a mildly elevated blood alcohol concentration. You make a note to yourself to ask her about this at her next visit with you.

When Isabella arrives at her psychiatric follow-up appointment, she appears even more visibly anxious than last time. She says that she tried to utilize some of the strategies that you shared with her previously, but she is only able to do them for a few minutes before her heart starts to pound and her mind starts to race. You ask her about her recent emergency department visit and again inquire if she has been using any substances as of late. She admits to not telling the full truth at her last appointment and says that she has been having several glasses of wine over the past week. She says that “alcohol has helped me feel calmer in the past. I know that it’s not good for you when you’re pregnant, but I don’t know what else I can do.” She has not vomited since her emergency department visit, is not tremulous or diaphoretic, and denies any perceptual disturbances. You take the time to inform her of the negative impacts of alcohol exposure during pregnancy on fetal development. She voices understanding but starts crying, saying “I’m a bad mom, I’m a bad mom, I’m so stressed out and I need help.” You provide empathetic support and tell her that you are there to help her. You recognize that racial disparities, lower socioeconomic status, and lack of support can

contribute to higher rates of substance use. You also inform her that anxiety and alcohol use can both lead to the other and that although alcohol may temporarily suppress anxiety while drinking, there can be a rebound effect that significantly increases anxiety afterwards.

She says that she is now interested in starting an anti-anxiety medication as well as psychotherapy. You recognize that many women who screen positive for mood issues during pregnancy do not receive adequate treatment and that barriers to culturally and linguistically appropriate care exist. You are eager to share treatment options with her. You provide her with the indications/risks/benefits/side effects of Zoloft, and she is agreeable with starting 25 mg daily and following up with you in another two weeks. You give her a list of Spanish-speaking therapists that she can contact to schedule a session. She tells you that she will stop drinking, which you encourage, but you also provide her with alcohol use resources, including AA, in Spanish.

Visit 3

At Isabella's next appointment with you, you notice that Isabella has a moderate-sized bruise on her right cheek and several smaller bruises on her right arm. She smiles when you reiterate that it may take several additional weeks for the anxiolytic effect to work. She says that she did not drink any alcohol since her last appointment with you because "I was scared to." She says that this past week, she was trying to get rid of the alcohol in her home as a way to prevent herself from drinking but that her partner intervened and got upset that she attempted to throw out the beer and wine. You ask her if she felt threatened or if anything physical took place. She shakes her head and says "no." You ask her if she feels safe at home, and she replies "yes, I do." You provide normalizing information about substance use coercion and strategize with her about how to navigate those situations. You understand that IPV against pregnant women is a major public health issue that can lead to adverse outcomes in both mother and fetus and that mental health and substance use coercion are common forms of abuse. You also recognize that women may be reluctant to share details about their relationships with healthcare providers. You remind her that you are there to help her and that if there is any time where she does not feel comfortable at home or fears for the safety of her baby or herself, that she can contact your office, go to the emergency department, or call 911. She nods in understanding. You ask her about her social support system; she says that she has an aunt that lives around 35 minutes away. She says that she could get extra help from her aunt if she needed it. You ask her if she would like any resources for domestic violence/abuse. (You let her know about the kinds of help DV programs can offer and ask if she'd like any resources

shelters. She politely declines. You discuss increasing her Zoloft to a more therapeutic dose of 50 mg daily, and she is in agreement. She says that she was able to schedule a therapy appointment with a Spanish-speaking therapist that will take place next week. Her next appointment with you will be in two weeks.

Visit 4

At Isabella's next appointment with you, she says that she has been feeling a bit better. She has been calling her aunt once a week to talk about her pregnancy, and she has found her aunt's words of support and knowledge of pregnancy to be comforting. She also says that she had a good first session with her therapist and that it is nice to have someone else to talk to about her life stressors. She denies any side effects with the increase in Zoloft. She still reports feeling anxious at times but has been better able to manage it with meditation and breathing techniques. When asked about how things are going with her partner, she replies "the same" and does not elaborate further. She says that she likes her OB-GYN and feels more prepared to handle her pregnancy. You and Isabella decide to keep her medication at its current dosage for now, and you encourage her to continue relying on her social support system, seeing her therapist, and using her coping skills. She thanks you for your help. You know that prenatal anxiety is a strong predictor of postpartum mood symptoms, so you recognize that after her delivery, she will need closer follow up with you, but for now, given her overall improvement, you will see her again in one month.

- Perinatal=Prenatal (conception to birth) + Postpartum (birth to 6 weeks post-delivery, though this phase can extend to one year).
- Prenatal anxiety is a strong predictor of postpartum depression and anxiety. ● Women of color are disproportionately less likely to receive accurate diagnosis and treatment for perinatal anxiety disorders than white women, a disparity exacerbated by racial discrimination and bias in the healthcare system. Additionally, cultural stigmas and taboos surrounding mental health in specific communities contribute to underreporting and underdiagnosis of perinatal anxiety disorders among women of color.
- While there are many evidence-based practices for perinatal mood and anxiety disorders, screening rates vary widely, and $\leq 25\%$ of those who screen positive for depression receive any treatment. Gaps in care loom largest for individuals who are Black/African American, Hispanic/Latina/o/e/x, American Indian or Alaska Native, or Native Hawaiian or other Pacific Islander (populations that have been marginalized).
- Adverse social determinants of health (SDOH) and disparities in screening or

treatment rates drive inequities in perinatal mental healthcare.

- While the maternal mortality rate for Latina women is lower than that for Black and Native American women, it is still higher than that for White women. Socioeconomic factors, immigration status, and access to culturally competent care play significant roles.
- Racial disparities, lower socioeconomic status, and lower social support contribute to higher rates of substance use disorders (SUD) which can lead to complications during pregnancy and birth.
- There is limited availability for specialized SUD programs designed for pregnant persons. Additionally, lack of access to SUD treatment disproportionately affects those with limited financial resources. Intra and extra community stigma related to substance use against minority perinatal patients can create barriers to care. Systemic racism also makes it more likely that a pregnant person of color would face harsher penalties for substance use compared with white women. This may be a barrier for pregnant persons of color to seek help. Other barriers include limited access to culturally and linguistically appropriate care and affordable access to care.
- Intimate partner violence against pregnant women is a public health concern that can contribute to adverse outcomes in both mother and fetus: physical trauma, maternal depression, anxiety, and substance use; low birth weight, preterm birth, and placental abruption.

Resources

King C, Laynor G, McNeely J, Fawole A, Lee M, Terplan M, Choi S. Strategies to improve delivery of equitable and evidence-informed care for pregnant and birthing people with a substance use disorder in acute care settings: A scoping review protocol. Plos one. 2024 Mar 18;19(3):e0300183.

Kushner, M. (2000). The relationship between anxiety disorders and alcohol use disorders A review of major perspectives and findings. Clinical Psychology Review, 20(2), 149–171. [https://doi.org/10.1016/s0272-7358\(99\)00027-6](https://doi.org/10.1016/s0272-7358(99)00027-6)

Navarrete L, Nieto L, Lara MA. Intimate partner violence and perinatal depression and anxiety: Social support as moderator among Mexican women. Sexual & Reproductive Healthcare. 2021 Feb 1;27:100569.

Perinatal Mental Health Toolkit. (2024). Psychiatry.org.
<https://www.psychiatry.org/psychiatrists/practice/professional-interests/women-s-mental-health/maternal-mental-health-toolkit>

Schiff DM, Li WZ, Work EC, Goullaud L, Vazquez J, Paulet T, Dorfman S, Selk S, Hoepfner BB, Wilens T, Bernstein JA. Multiple marginalized identities: A qualitative exploration of intersectional perinatal experiences of birthing people of color with substance use disorder in Massachusetts. *Journal of Substance Use and Addiction Treatment*. 2024 May 23:209346.

Sufredini F, Catling , Zugai J, Chang S. The effects of social support on depression and anxiety in the perinatal period: A mixed-methods systematic review. *Journal of Affective Disorders*. 2022 Dec 15;319:119-41.

VIGNETTE #2

Visit 1 - First trimester

Janelle is a 28 year old female with a past psychiatric diagnosis of schizoaffective disorder, bipolar type who presents to your office after her sister recommended that she see a psychiatrist. She scored a 3 on the Edinburgh Postnatal Depression Scale that she completed before her appointment. She just found out that she is pregnant second child and her sister wants to make sure that she's healthy during this pregnancy. She had been prescribed psychiatric medication in the past but she discontinued her medications because she doesn't want to "hurt the baby" with problems that are "all in her head". She said that she heard from her family that medications can cause birth defects.

You gather additional information about her past psychiatric history. She explains that she was diagnosed with schizoaffective disorder when she was 19. Review of symptoms during that time is significant for irritability. She also explained, "the doctors said I was paranoid." You consider the roles of bias in psychotic spectrum diagnoses and cultural paranoia. You explore her past psychiatric history further and realize that she has a history of sexual trauma and that her symptoms of irritability, wariness towards males in a hospital setting (interpreted as paranoia), avoidance, flashbacks, and poor sleep most closely fit with PTSD as a result of her sexual trauma. Per review of symptoms, she has not had any symptoms consistent with a manic episode or true psychosis.

You consider the disparities in maternal mortality and you recommend that she follow up with her OB visits in addition to seeing you for appointments.

You explore her symptoms after her first pregnancy and she stated that she had increased irritability, anhedonia, difficulty connecting with her baby, and suicidal thoughts after the birth of her first child. She also had unwanted thoughts about harming her baby and they scared her. She reports that her strong faith Christian prevented her from acting on these homicidal thoughts. She goes to church weekly and feels community at her church but is worried about being judged for her thoughts. She reported that she was worried disclosing her unwanted homicidal thoughts towards her child would result in her child getting taken away by DSS. You acknowledge her fears that there is still a stigma towards mental illness and ask further questions about her understanding of why she's experiencing these symptoms and how others within her community interpret similar symptoms. You use elements of the cultural formulation interview ([APA DSM5 Cultural-Formulation-Interview.pdf](#)). She explains that she worries that it's her fault that she wasn't able to connect the way she wanted to her baby. She states that her pastor is very compassionate and welcoming. She has never talked to her pastor about her symptoms but she remembers her pastor leading a prayer for a community member who was in the hospital for their mental health.

You ask whether she would consider including her pastor in her care plan moving forward and she agrees. She ultimately declines medication for her history of PTSD and postpartum depressive symptoms at this time. You review the symptoms of depression with her and discuss warning signs that should prompt her to return for an appointment and symptoms that are considered emergencies. You also review the risks of untreated depression compared with the risks of taking medication to treat depression during pregnancy. You are respectful of this decision and explain that you are here if she changes her mind in the future.

Follow up visit: Postpartum

The patient has had worsening depressive symptoms including anhedonia, irritability, and suicidal thoughts. She scored a 13 on the Edinburgh Postnatal Depression Scale. She states that the symptoms began during her pregnancy and have not gotten better. The symptoms became so bad that she wasn't able to continue working. She is very withdrawn and flat on exam.

You consider that her symptoms impacting her work may have affected her ability to access safe and stable housing, food, and other necessities. You do a brief screen to better understand these issues. She explains that she was able to get additional support from her church community for now but is worried about the future. Your clinic has a social worker because of how closely intertwined mental health is with social

determinants of health. You connect her with the clinic social worker who can help explain the resources available in your area. If your clinic doesn't have a social worker, your clinic leadership compiled a list of local resources where the patient can get help.

You discuss treatment options with her which include medication and therapy. She explains that she's willing to do therapy as long as the provider is respectful of her beliefs and integrates those beliefs into the treatment plan. You refer her to someone who practices culturally informed therapy.

- Maternal mortality refers to deaths due to complications from pregnancy or up to a year following childbirth. Black women are disproportionately affected, with maternal mortality rates approximately three times higher than those of white women. This disparity persists regardless of income or education level, highlighting systemic issues in healthcare access and quality.
- SDOH are the social circumstances in which people are born, work, live, and age and the wider set of forces shaping daily life. SDOH influence 80% of health outcomes as well as patient engagement with the healthcare system. Addressing patients' adverse SDOH is critical to improving health equity and is essential for high-quality healthcare. Professional organizations and policy makers recommend screening for adverse SDOH at healthcare visits. Integrating social services into mental healthcare is critical for improving health equity and patient experience.
- While effective treatments are available, perinatal mental health care often falls short of patients' needs and of recommended standards. Barriers persist across the mental healthcare pathway (screening, assessment, and treatment until symptom remission), including stigma, fear regarding treatment, and a dearth of community-based mental health supports. Providers and patients report that barriers are amplified by 1) gaps in provider training, 2) lack of established treatment workflows, 3) lack of access to mental healthcare resources, referrals, and consultation, and 4) microaggressions, implicit racial biases, and discrimination. Gaps and inequities persist across the perinatal mental healthcare pathway. Thus, the US lags behind other high-income countries.
- Professional organizations and policymakers recommend screening for adverse SDOH at healthcare visits. Integrating social services into mental healthcare is critical for improving health equity and patient experience.
- Black women are disproportionately affected, with maternal mortality rates approximately three times higher than those of white women. This disparity persists regardless of income or education level, highlighting systemic issues in healthcare access and quality.

- Maternal mortality refers to deaths due to complications from pregnancy or up to a year following childbirth. In the United States, a number of disparities affect the maternal mortality rate including access to quality care, race/ethnicity, low-income, lack of education, and language barriers
- Implicit biases in healthcare, discriminatory practices, and historical mistrust contribute to these disparities. Chronic conditions like hypertension, diabetes, and obesity, which are more prevalent in some racial groups, increase maternal mortality risk. Poverty, lack of education, and inadequate insurance coverage disproportionately affect minorities, impacting maternal health outcomes.
- Efforts to reduce these disparities include improving healthcare access, addressing social determinants of health, and increasing awareness and training among healthcare providers to deliver culturally competent care are all ways to reduce these disparities.
- Black and Hispanic women often face delayed diagnoses or are misdiagnosed with other conditions such as depression or schizophrenia, rather than postpartum psychosis. Mental health symptoms in women of color may be dismissed or underestimated, leading to a failure in timely recognition of the severity of their condition.

Resources

American Psychiatric Association. (2013). Cultural Formulation Interview (CFI).

American Psychiatric Association.

https://www.psychiatry.org/File%20Library/Psychiatrists/Practice/DSM/APA_DSM5_Cultural-Formulation-Interview.pdf

APA Official Actions Position Statement on Screening and Treatment of Mood and Anxiety Disorders During Pregnancy and Postpartum.

<https://www.psychiatry.org/getattachment/bf2c29a1-98f5-4c8f-9e4a-4119fa64bcca/Position-Pregnancy-Postpartum-Mood-Anxiety-Disorders.pdf>

Faber, S. C., Khanna Roy, A., Michaels, T. I., & Williams, M. T. (2023). The weaponization of medicine: Early psychosis in the Black community and the need for racially informed mental healthcare. *Frontiers in Psychiatry*, 14(1098292).

<https://doi.org/10.3389/fpsy.2023.1098292>

Rainosek, C. (2023, August 24). How to Show Compassion and Sensitivity While

Screening for Social Needs. Salud America.

<https://salud-america.org/how-to-show-compassion-and-sensitivity-while-screening-for-social-needs/>

Weisman de Mamani, A., Altamirano, O., Lopez, D., McLaughlin, M. M., Maura, J., Martinez de Andino, A., Ahmad, S. S., Hafner, L., & Lund, S. G. (2023). Culturally informed therapy: An intervention that addresses the psychological needs of religious individuals of diverse identities. In P. S. Richards, G. E. K. Allen, & D. K. Judd (Eds.), *Handbook of spiritually integrated psychotherapies* (pp. 33–55). American Psychological Association. <https://doi.org/10.1037/0000338-002>